

SHILAP Revista de Lepidopterología

ISSN: 0300-5267 avives@eresmas.net

Sociedad Hispano-Luso-Americana de Lepidopterología España

Asselbergs, J. E. F.; Seguna, A.; Sammut, P.
Recent records of Pyraloidea species new to Malta, including two species new to the European fauna (Lepidoptera: Pyraloidea)

SHILAP Revista de Lepidopterología, vol. 36, núm. 144, diciembre, 2008, pp. 465-471

Sociedad Hispano-Luso-Americana de Lepidopterología

Madrid, España

Available in: http://www.redalyc.org/articulo.oa?id=45511220008



Complete issue

More information about this article

Journal's homepage in redalyc.org



Recent records of Pyraloidea species new to Malta, including two species new to the European fauna (Lepidoptera: Pyraloidea)

CODEN: SRLPEF

J. E. F. Asselbergs, A. Seguna & P. Sammut

Abstract

Acrobasis obliqua clusinella Zeller, 1848, Acrobasis centunculella (Mann, 1859), Psorosa dahliella (Treitschke, 1832), Psorosa mediterranella Amsel, 1953, Euzopherodes lutisignella (Mann, 1869), Delplanqueia inscriptella (Duponchel, 1836), Isauria dilucidella (Duponchel, 1836), Euchromius gozmanyi Błeszyński, 1961, Udea numeralis (Hübner, 1796), Ancylosis (Heterographis) costistrigella (Ragonot, 1890) and Caina deletella Ragonot, 1893 are new to the Maltese Islands. The latter two species are also the first records for the European fauna. Where known, data on the biology, the first stages and the distribution of the species is given. Additional notes on some other pyraloid species from Malta are also given.

KEY WORDS: Lepidoptera, Pyraloidea, new records, European fauna, Malta.

Recientes citas de especies nuevas de Pyraloidea para Malta, incluyendo dos especies nuevas para la fauna europea (Lepidoptera: Pyraloidea)

Resumen

Acrobasis obliqua clusinella Zeller, 1848, Acrobasis centunculella (Mann, 1859), Psorosa dahliella (Treitschke, 1832), Psorosa mediterranella Amsel, 1953, Euzopherodes lutisignella (Mann, 1869), Delplanqueia inscriptella (Duponchel, 1836), Isauria dilucidella (Duponchel, 1836), Euchromius gozmanyi Błeszyński, 1961, Udea numeralis (Hübner, 1796), Ancylosis (Heterographis) costistrigella (Ragonot, 1890) y Caina deletella Ragonot, 1893, son nuevas para Malta. Las dos últimas especies también son primeras citas para la fauna europea. Se dan, donde se conocen, datos sobre su biología, los primeros estados y la distribución de las especies. También se dan notas de algunas otras especies de pirálidos de Malta.

PALABRAS CLAVE: Lepidoptera, Pyraloidea, nuevas citas, fauna europea, Malta.

Introduction

So far, about 16,000 Pyraloidea species have been described (HEPPNER, 1991). It is estimated that many more are still awaiting description. Also, from many species the early stages, the biology and the distribution are insufficiently known or not known at all. In this paper eight Pyraloidea species new to Malta are mentioned, including two which are new to the European fauna. At present the Pyraloidea-fauna of Malta comprises 134 species, which number is composed as follows: Pyralidae are represented by Galleriinae (5 species), Pyralinae (12 species) and Phycitinae (68 species) species; Crambidae are represented by Scopariinae, Cybalomiinae and Acentropinae (1 species each), Glaphyriinae (2 species), Odontiinae and Evergestinae (3 species each), Crambinae (9 species), Spilomelinae (13 species) and Pyraustinae (16 species) (SAMMUT, 1984, 2000).

Material and methods

In this article, unless otherwise stated, all material has been determined by the first author; the material studied is in the collections of the second and the third author and has been collected at mixed UV/MV light. All material studied is in the collection of the second two authors.

PYRALIDAE Phycitinae

Acrobasis obliqua clusinella Zeller, 1848 (Fig. 1)

[Acrobasis clusinella Zeller, 1848. Isis von Oken, 1848: 610]

Material examined: MALTA, 1 ex., Bingemma, 30-IV-1999, A. Seguna leg.; 1 ex., Mellieha, il-Kortin, 12-IV-2006, H. Borg-Barthet leg. [in coll. Sammut]; 1 ex., Naxxar, Tas-Saghjtar, 29-IV-2006, A. Seguna leg.; 1 $\,^{\circ}$, Rabat, Dwejra, 25-IV-2007, A. Seguna leg.; 1 ex., Rabat, Dwejra, 30-IV-2007, A. Seguna leg.

Biology: The larva of the nominotypical form is known to feed on *Cistus* spp. (SPULER, 1910). Adults from V-VIII.

Distribution: Known mainly from S. and S.E. Europe, including the Mediterranean islands of Corsica, Sardinia and Sicily (NUSS *et al.*, 2004).

Acrobasis centunculella (Mann, 1859) (Fig. 2)

[Myelois centunculella Mann, 1859. Wien. ent. Monatschr., 3: 165]

Material examined: MALTA, 2 ♀ ♀, Rabat, Dwejra, 25-IV-2007, A. Seguna leg.

Biology and first stages undescribed. Adults from IV-VIII.

Distribution: Known from France, Spain, Italy and Sicily, former Yugoslavia, Greece, and Turkey (NUSS et al., 2004)

Psorosa dahliella (Treitschke, 1832) (Fig. 3)

[Phycis dahliella Treitschke, 1832. Schmett. Europa, 9(1):198]

Material examined: MALTA, 1 ♂, Naxxar, 11-VII-2001, A. Seguna leg.

Biology and first stages unknown. Adults: from VI-X.

Distribution: Known from S. and SE. Europe, Slovakia, Corsica, Sardinia and Sicily (NUSS et al., 2004).

Psorosa mediterranella Amsel, 1953 (Fig. 4)

[Psorosa mediterranella Amsel, 1953. Ark. Zool., [2]6: 279, Figs. 27a, 29]

Material examined: MALTA, 13, Rabat, Dwejra, 28-VI-2004, [gen. prep. GP 5796 Ass.] A. Seguna leg.; 13, Naxxar, Tas-Sghajtar, 20-V-2006, A. Seguna leg.

Biology: early stages and larval host plant unknown. Adults from VII-IX.

Distribution: France, Spain, Portugal, Sicily, Italy and former Yugoslavia ((NUSS et al., 2004).

Ancylosis (Heterographis) costistrigella (Ragonot, 1890) (Fig. 5)

[Heterographis costistrigella Ragonot, 1890. Bull. Soc. ent. Fr., 1890: 111]

Material examined: MALTA, 18, Naxxar, 22-V-2006, A. Seguna leg.

Biology: Larva feeds in silken tubes at ground level, mining leaves of *Salvia aegyptiaca* L. mining the leaves (CHRÉTIEN, 1917). Locally, the larval host plant is not yet known.

Distribution: Known from Morocco, Algeria, Tunisia, Libya, Bahrain, Palestine, Sudan, Iran, and Afghanistan (ROESLER, 1973).

Euzopherodes lutisignella (Mann, 1869) (Fig. 6)

[Myelois lutisignella Mann, 1869. Verh. zool.-bot. Ges. Wien, 19: 381]

Material examined: MALTA, $1\$ \$\ Siggiewi, Ghar Lapsi, 3-VIII-2001, [gen. prep. GP 5818 Ass.], P. Sammut leg.; $1\$ \$\ Naxxar, 8-VI-2003, A. Seguna leg.; $1\$ \$\ Gharghur, 10-IV-2007, A. Seguna leg.; $1\$ \$\ Rabat, Dwejra, 25-IV-2007, A. Seguna leg.; $1\$ \$\ Rabat, Dwejra, 30-IV-2007, [gen. prep. GP. 5791 Ass.], A. Seguna leg.; $1\$ \$\ Naxxar, 20-VI-2007, A. Seguna leg.

Biology and first stages: Larva on *Phlomis lychnitis* L. (HAMPSON, 1901). Adults: from III-VIII (based on label information from specimens in the collection of the first author).

Distribution: In Europe, the nominotypical subspecies *lutisignella*, is known from Sicily, Romania, former Yugoslavia, Albania, Bulgaria, Greece and Crete (NUSS *et al.*, 2004). In N. Africa it has been recorded from Cyrenaica (ROESLER, 1973).

Delplanqueia inscriptella (Duponchel, 1836) (Fig. 7)

[Phycis inscriptella Duponchel, 1836. In Godart & Duponchel, Hist. nat. Lépid. Fr. 10(7): 202, pl. 279]

Material examined: MALTA, 1 ♀, Naxxar, 16-VI-2002, [gen. prep. GP 5789 Ass.], A. Seguna leg. Biology: Larva on *Thymus* sp. (LERAUT, 2001).

Distribution: In Europe it is known from Spain and France (NUSS et al., 2004; LERAUT, 2001)

Isauria dilucidella (Duponchel, 1836) (Fig. 8)

[Phycis dilucidella Duponchel, 1836. In Godart & Duponchel, Hist. nat. Lépid. Fr., 10:184, pl. 278, fig.7]

Material examined: MALTA, 1 ex., Naxxar, 6-V-1999, A. Seguna leg.; 1 $\,^{\circ}$, Rabat, 6-X-2001, P. Sammut leg.; 1 $\,^{\circ}$, Rabat, 13-IX-2002, P. Sammut leg.; 1 $\,^{\circ}$, Dingli, Dingli Cliffs, 10-VI-2003, [gen. prep. GP. 5806 Ass.], A. Seguna leg.; 1 ex., Dingli, Dingli Cliffs, 4-X-2004, A. Seguna leg.; 1 ex., Rabat, Dwejra, 9-VI-2007, A. Seguna leg.

Biology: Larva is reddish-brown, yellowish ventrally with flown-out, slightly darker side lines (HASENFUSS, 1960). Head yellow (HANNEMANN, 1964). Larva can be found from July, and, after hibernation till June of the following year, in a silken tube perpendicular in the soil, feeding on *Lotus corniculatus* L., and *Astralagus monspessulanus* L. (LHOMME, 1935). Adults: are on the wing from April till September.

Distribution: In Europe, the species is found in Spain, France, Switzerland, Austria, Germany, Poland, Czech Rep., Slovakia, Italy, Sicily, Sardinia, Hungary, Romania, former Yugoslavia, Bulgaria, Albania, Greece, and European Russia (NUSS *et al.*, 2004). From N. Africa it is known from Algeria (CHRETIEN, 1917). It is also known from Turkey, Syria, Lebanon, Palestine, Iran, Iraq, Afghanistan, Pakistan, Turkmenistan and Mongolia (ROESLER, 1993).

Caina deletella Ragonot, 1893 (Fig. 9)

[Caina deletella Ragonot, 1893. In Romanoff, Mém. Lép., 7: 466, pl.18, fig.11]

Material examined: MALTA, 1 \, Rabat, 28-III-2007, [GP. 5814 Ass.], P. Sammut leg.

Biology and first stages unknown. Adults are on the wing from the end of March (this paper) till June and again from September till November (RAGONOT, 1893), probably in two generations.

Distribution: Known from the Persian Gulf (Fao) and from India (Poona) (RAGONOT, 1893). This species is new also to the fauna of Europe.

CRAMBIDAE Crambinae

Euchromius gozmanyi Błeszyński, 1961 (Fig.10)

[Euchromius gozmanyi Błeszyński, 1961. Acta ent. Mus. natn. Pragae, 34: 462, figs. 7, 17]

Material examined: GOZO, 1 ex., Kercem, San Rafflu, 15-VI-2001, P. Sammut leg.; 9 exx., Ramla l-Hamra, 19-VI-1998, A. Seguna leg.; 5 exx., Ramla l-Hamra, sand dunes, 9-VIII-2002, P. Sammut & A. Seguna leg.; 39 exx., Marsalforn, Qbajjar, 25-V-2005, P. Sammut & A. Seguna leg.; 25 exx., same

data, 8-VI-2006; GP 5853 Ass., male and 5854 Ass., male, P. Sammut & A. Seguna leg.; 2 exx., Xaghra, 21-23-VII-2005, A. Catania leg.; 5 exx., same data but 11-VII-2005, A. Catania leg.; MALTA, 1 ex., Cirkewwa, Paradise Bay, 18-V-1997, A. Seguna leg.; 3 exx., same data but 10-VI-1998, A. Seguna leg.; 1 ex., Mellieha, Marfa Ridge, Rdum tal-Madonna, 8-VIII-2007, P. Sammut leg.; 1 ex., Mistra, 16-IX-1995, A. Catania leg.; 1 ex., Naxxar, 2-V-1997, A. Seguna leg.

Biology and early stages undescribed (BŁESZYŃSKI, 1965). On the Maltese islands, the species appears to be either two brooded from May to June and again during August and September; elsewhere it produces one continuous long brood.

Distribution: In Europe the species has been recorded from The Iberian peninsula, France, Sardinia and Sicily (BŁESZYŃSKI 1965; NUSS *et al.*, 2004). From the North African coast, it has been recorded from Ain Draham in Tunisia and from Algeria (BŁESZYŃSKI, 1965). According to RUNGS (1980) it is also known from Morocco.

CRAMBIDAE Pyraustinae

Udea numeralis (Hübner, 1796) (Fig. 11)

[Pyralis numeralis Hübner, 1796. Samml. europ. Schmett. Pyr.: pl. 14, fig. 89]

Material examined: MALTA, 1 ex., Dingli, Dingli Cliffs,18-IV-2004, A.Seguna leg.; 1 ex., Dingli, Dingli Cliffs, 5-X-2005, P. Sammut leg.; 1 ex., Mellieha, il-Kortin, 12-VI-2005, H. Borg-Barthet leg.; 1 ex., Mellieha, Mgiebah, 15-X-2004, A. Seguna leg.; 1 ex., Naxxar, 5-V-2005, A. Seguna leg.; 1 ex., Rabat, Dwejra, 23-X-2003, A. Seguna leg.; 1 ex., Rabat, Dwejra, 20-X-2004, P. Sammut leg.; 1 ex., Rabat, Dwejra, 21-X-2004, A. Seguna leg.; 1 ex., Rabat, Dwejra, 30-X-2004, A. Seguna leg.; 1 ex., Rabat, Dwejra, 27-V-2006, A. Seguna leg.; 1 d, Rabat, Dwejra, 28-X-2006, A. Seguna leg.; 1 ex., Rabat, Dwejra, 29-X-2007, A. Seguna leg.; 1 ex., Rabat, Dwejra, 12-V-2007, A. Seguna leg.; 1 ex., Rabat, Dwejra, 12-V-2007, A. Seguna leg.; 1 ex., Rabat, Dwejra, 12-V-2007, A. Seguna leg.

Biology: Larva on Brassicaceae (SPULER, 1910; SLAMKA, 1997).

Distribution: Southern European species known from France, Spain, Portugal, Corsica, Sardinia, Sicily, Italy, former Yugoslavia, Greece and Turkey.

Conclusion

Besides the above mentioned species which are all new to the pyraloid fauna of the Maltese archipelago, other species, such as *Denticera divisella* (Duponchel, 1842), *Neurotomia coenulentella* (Zeller, 1846), *Epischnia peroni* Agenjo, 1948, *Bradyrrhoa cantenerella* (Duponchel, 1837), *Postemmalocera palaearctella* (Turati, 1917), *Tegostoma comparalis* (Hübner, 1796), and *Sitochroa palealis* ([Denis & Schiffermüller], 1775), formerly considered as rare species have been recorded frequently from different stations on the Maltese Islands. Numerous examples of *Ancylolomia tripolitella* Rebel, 1909 have been recorded at light from l-Irdum tal-Madonna at Marfa Ridge.

Finally *Ancylolomia minutella* Turati, [1926], recorded by SAMMUT (1984, 2000) is not this species but *Ancylolomia tentaculella* (Zeller, 1847).

Acknowledgements

We would like to thank Dr. František Slamka (Bratislava, Slovakia) for pointing out to us the correct identity of *Ancylolomia minutella*, Mr. Aldo Catania for providing data on *Euchromius gozmanyi* in his collection, and Dr. Antonio Vives for providing the Spanish translation of the abstract.

BIBLIOGRAFÍA


AMSEL, H. G., 1953.– Die Microlepidopteren der Brandt'schen Iran-Ausbeute 4 Teil.– *Arkiv Zool.*, **6**(26): 255-326, 33 pls.

- BŁESZYŃSKI, J., 1965.– Crambinae. *In H. G. AMSEL*, F. GREGOR & H. REISSER (eds.).– *Microlepidoptera Palaearctica*, 1: 553 pp., 133 pls. Georg Fromme, Wien.
- BORG, P., 1932.- The Lepidoptera of the Maltese Islands: v + 25 pp. Government Printing Press, Malta.
- CHRÉTIEN, P., 1917. Contribution à la connaissance des Lépidoptères du Nord de l'Afrique. Notes biologiques et critiques. *Annls Soc. ent. Fr.*, **85** (1916): 369-503.
- FILIPJEV, N. N., 1931.— Lepidopterologische Notizen XIII. Ein neuer Weintrauben-schädling aus der Gattung Ephestia Guenée.— Dt. ent. Z. Iris, 45: 70-73.
- HAMPSON, G. F., 1901.– Monographie des Phycitinae et des Galleriinae II. *In* N. M. ROMANOFF.– *Mémoires sur les Lépidoptères*, **8**: 602 pp., 57 pls. St. Pétersbourg.
- HANNEMANN, H. J., 1964.— Kleinschmetterlinge oder Microlepidoptera II. Die Wickler (s. l.). (Cochylidae und Carposinidae). Die Zünslerartigen (Pyraloidea).— *Tierwelt Dtl.*, **50**: VIII + 401 pp., 22 pls.
- HASENFUSS, I., 1960.– Die Larvalsystematik der Zünsler (Pyralidae).– Abh. Larvalsyst. Insekten, 5: 10-263, 219 figs., 2 pls.
- HEPPNER, J. B., 1991. Faunal regions and the diversity of Lepidoptera. Tropical Lepid., 2, suppl., 1: 1-85.
- LERAUT, P., 2001. Contribution à l'étude des phycites palaearctiques (Lepidoptera, Pyralidae, Phycitinae). Revue fr. Ent. (N. S.), 23 (2):129-141.
- LERAUT, P., 2002. Contribution à l'étude des Phycitinae (Lépidoptera, Pyralidae).— Nouv. Revue Ent. (N. S.), 19 (2): 141-177.
- LHOMME, L., 1935-[1946].— Catalogue des Lépidoptères de France et de Belgique, 2(1): 487 pp. La Carriol, Par Douelle (Lot).
- NUSS, M., SPEIDEL, W. & SEGERER, A., 2004. Fauna Europaea: Pyralidae. In O. KARSHOLT & J. RAZOWSKI (eds). Fauna Europaea version 1.1, http://www.faunaeur.org
- RAGONOT, E. L., 1893. Monographie des Phycitinae et des Galleriinae I. In N. M. ROMANOFF. Mémoires sur les Lépidoptères, 7: 658 pp., 28 pl.
- ROESLER, R. U., 1973. Trifine Acrobasiina 1. Teilband der Phycitinae. *In* H. G. AMSEL, F. GREGOR & H. REISSER (eds). *Microlepidoptera Palaearctica*, 4: 752 pp., 170 pls. Georg Fromme & Co., Wien.
- RUNGS, Ch., 1980. Catalogue raisonné des Lépidoptères du Maroc. Inventaire faunistique et observations écologiques. *Trav. Inst. scient. chérif. Sér. Zoologique*, 1: 1-222, 2 cartes (1979).
- SAMMUT, P., 1984. A Systematic and Synonymic List of the Lepidoptera of the Maltese Islands.— *Neue Ent. Nachr.*, 13: 1-124.
- SLAMKA, F., 1997. Die Zünslerartigen (Pyraloidea) Mitteleuropas: 112 pp., 55 + 13 pls. F. Slamka, Bratislava.
- SPEIDEL, W., 1996.– Pyralidae. In O. KARSHOLT & J. RAZOWSKI (eds).– Checklist of the Lepidoptera of Europe: 380 pp. Apollo Books, Stenstrup.
- SPULER, A., 1910. Die Schmetterlinge Europas- Kleinschmetterlinge: pp.188-523, Tafeln 81-91. Stuttgart.


J. E. F. A. Neerland 20 NL-4614 GD Bergen op Zoom HOLANDA / THE NETHERLANDS E-mail: jef.asselbergs@hetnet.nl A. S. 68, 'Redeemer', Triq L-Emigranti Naxxar NXR 3200 MALTA / MALTA E-mail: seguna@onvol.net

P. S. 137, 'Fawkner/2' Dingli Road Rabat RBT 9023 MALTA / MALTA E-mail: farfett@onvol.net

(Recibido para publicación / Received for publication 20-VI-2008) (Revisado y aceptado / Revised and accepted 11-VII-2008)



Figs. 1-6.– 1. Acrobasis obliqua clusinella Z. Malta, Rabat, Dwejra, 25-IV-2007. **2.** Acrobasis centuculella (Mn.). Malta, Rabat, 25-IV-2007. **3.** Psorosa dahliella (Tr.). Malta, Naxxar, 11-VII-2001. **4.** Psorosa mediterranella Ams. Malta, Rabat, Dwejra, 28-VI-2004. **5.** Ancylosis costistrigella (Rag.). Malta, Naxxar, 22-V-2006. **6.** Euzopherodes lutisignella (Rag.). Malta, Rabat, Dwejra, 30-VI-2007.



Figs. 7-11.— 7. Delplanqueia inscriptella (Dup.). Malta, Naxxar, 16-VI-2002. 8. Isauria dilucidella (Dup.). Malta, Dingli, Dingli Cliffs, 10-VI-2003. 9. Caina deletella (Dup.). Malta, Rabat, 28-III-2007. 10. Euchromius gozmayi Błesz. Malta, Gozo, Kercem, San Rafflu, 19-VI-1998. 11. Udea numeralis (Hb.). Malta, Dwejra, 28-X-2006.